

RURAL AND RACIAL DISPARITIES IN COLORECTAL CANCER INCIDENCE AND MORTALITY IN SOUTH CAROLINA, 1996 - 2016

R Ranganathan, SA Adams, WE Zahnd, P Hung, MJ Brown, JM Eberth


Rural and Minority Health Research Center, Department of Epidemiology & Biostatistics, College of Nursing & Department of Health Services Policy and Management, University of South Carolina, USA



PURPOSE

- Colorectal cancer (CRC) is the third leading cause of cancer mortality in the United States and South Carolina (SC) and its age-adjusted incidence and mortality are highest among Black population. [1]
- With a high proportion of Black (27.9%) and rural residents (33.7%), it is important to describe the burden of CRC among these underserved populations in SC [2] as reducing the incidence of CRC remains one of the Healthy people 2020's objective priorities.

Maps of CRC incidence, 2010 – 2015, A) County level MIRs Vs SC MIR; B) County MIRs Vs U.S MIR; C) County level incidence rates by meeting the Healthy People 2020 goals



RESULTS

- Areas with high MIRs tended to be rural counties.
- Rural residents had higher proportions of distant stage CRC compared to urban residents, and Black population had higher proportions of distant stage CRC compared to Whites (26.3% vs. 22.7% and 29.3% vs. 23.7%, respectively; p-value < 0.05).
- From 1996-2016, Black and White urban dwelling-residents experienced a significant decline in incidence. Urban Whites, urban Blacks, and rural Whites experienced significant declines in mortality (AAPC= -2.6%UW vs. -2.4%UB vs. -1.6% RW vs. -0.9%RB, respectively).

METHODS

- CRC incidence, staging and mortality data were obtained from the SC Central Cancer Registry.
- Urban-rural status was determined Urban Influence Codes (1-2=urban; 3-12=rural).
- Mortality-to-incidence ratios (MIRs) for each SC county (2012-2016 period) were calculated and mapped using ArcGIS Version 10.5.1.
- Chi-square tests were calculated to examine differences in CRC stage by urban-rural status and race.
- Annual percent change (APC) and annual average percent change (AAPC) were calculated to examine trends in incidence and mortality rates across urban-rural and racial subgroups between 1996 and 2016.

CONCLUSIONS

Despite improvements in CRC screening in recent years, focused evidenced-based interventions for lowering incidence and mortality through improving early detection and treatment among rural and Black populations in SC are necessary.


REFERENCES

[1] Society A. *American Cancer Society. Colorectal Cancer Facts & Figures 2017-2019.*; 2019. <https://www.cancer.org/research/cancer-facts-statistics/all-cancer-facts-figures/cancer-facts-figures-2017.html>
 [2] Health and Demographics Division. South Carolina Statistical Abstract. Accessed June 12, 2020. <http://abstract.sc.gov/index.html>

ACKNOWLEDGEMENT

This study was supported by the Federal Office of Rural Health Policy (FORHP), Health Resources and Services Administration (HRSA), US Department of Health and Human Services (HHS) under cooperative agreement no. U1CRH30539. The information, conclusions, and opinions expressed in this article are those of the authors and no endorsement by FORHP, HRSA, or HHS is intended or should be inferred.

Age adjusted CRC Incidence & Mortality Rates, 1996 - 2016



AAPC=average annual percent change; *=statistically significant trend at p<0.05