

# The Association of a Rural Hospital Closure with 30-Day Post Hospital Discharge Mortality from Selected Conditions

Melinda A. Merrell, MPH<sup>1,2</sup>, Janice Probst, PhD<sup>1</sup>,  
Kevin Bennett, PhD<sup>3</sup>, Bankole Olatosi, PhD<sup>1</sup>, Jeffrey Hall, MD<sup>3</sup>

University of South Carolina<sup>1</sup>, South Carolina Office of Rural Health<sup>2</sup>, University of South Carolina School of Medicine<sup>3</sup>

## BACKGROUND AND PURPOSE

- During the last decade, over 100 rural hospitals have closed in the United States<sup>1</sup>.
- Concurrently, rural populations, compared to urban, are experiencing disparate trends in all-cause, cause-specific, premature, and in-hospital mortality rates<sup>2-5</sup>.
- It is unknown what impact, if any, rural hospital closures will have on these mortality trends.
- 30-day post hospital discharge mortality is a CMS quality indicator that allows for the standardization of mortality outcomes across time and facility<sup>6</sup>.

### Purpose:

Our study examines, in one state, the association of rural hospital closure with 30-day post hospital discharge mortality for selected conditions impacted most by timely access to health care services.

## METHODS

- Data were obtained for one Southeastern U.S. state from its all-payer claims database. 30-day post hospital discharge mortality was linked to individual encounters via the state's Vital Records department.
- Inpatient and emergency department (ED) patient encounters from study counties with ICD-9-CM diagnosis codes for acute myocardial infarction (AMI), stroke, sepsis, and trauma were included in the study.
- Study counties were chosen based on their rural community hospital status over the 60-month study period: **closure** occurred, **open** hospital, and **no** hospital. The closure county was identified first and statistical matching was used to identify four additional counties (2 with open hospitals, 2 with no hospital).
- Demographic characteristics of patients included age, gender, race, payment source, Charlson Index (co-morbidity score), and ED utilization during encounter.


## RESULTS

### Pre-closure time period characteristics of study sample (n=3,029):

- 30-day post hospital discharge mortality = 7.8% (n=236)
- Mean age = 62.0
- Gender = 48.9% female
- Race = 52.6% white
- Emergency Care Sensitive Condition Diagnoses:
  - AMI = 20.3%
  - Stroke = 24.2%
  - Sepsis = 39.2%
  - Trauma = 16.3%
- Significant differences by county hospital closure status were found for race, payment source, Charlson Index score, ED utilization, and sepsis and trauma diagnoses; adjusted in multivariable analysis.

### Patient characteristics associated with 30-day post hospital discharge mortality in pre-closure time period:

- Age (each additional year)
- Gender (female v. male)
- Medicare & Self-Pay payment sources (v. total)
- Charlson Index score (each incremental increase)
- ED utilization (yes v. no)
- Sepsis & Trauma Diagnoses (yes v. no)



### Key Findings

- *Logistic regression analyses showed no significant difference in 30-day post hospital discharge mortality rates between pre and post closure time periods in both unadjusted and adjusted models.*
- *However, for encounters from the closure county, the rate of change slows significantly in the post-closure time period in both unadjusted and adjusted models.*

## DISCUSSION

- In our study, 30-day post hospital discharge mortality rates from selected conditions were associated with patient's county of residence hospital closure status.
- It is unknown whether this finding represents improved access for these conditions or increased pre-hospital / in-hospital mortality for residents of counties with these conditions experiencing rural hospital closure.
- Additional analyses are needed. This study only represents one state/one closure in the Southeastern U.S. The explicit impact on vulnerable populations should also be explored further.

## ACKNOWLEDGEMENTS & REFERENCES

1. Cecil G. Sheps Center for Health Services Research. (2019). 104 Rural Hospital Closures: January 2010 – Present. Retrieved April 25, 2019, from <http://www.shepscenter.unc.edu/programs-projects/rural-health/rural-hospital-closures/>
2. Hoffman, A., & Holmes, M. (2017). Regional Differences in Rural and Urban Mortality Trends. Findings Brief, NC Rural Health Research Program, August 2017.
3. James, W. L. (2014). All rural places are not created equal: Revisiting the rural mortality penalty in the United States. *American Journal of Public Health*, 104(11), 2122–2129.
4. Mansfield, C. J., Wilson, J. L., Kobrinski, E. J., & Mitchell, J. (1999). Premature mortality in the United States: The roles of geographic area, socioeconomic status, household type, and availability of medical care. *American Journal of Public Health*, 89(6), 893–898.
5. Villapiano, N., Iwashyna, T. J., & Davis, M. M. (2017). Worsening Rural-Urban Gap in Hospital Mortality. *The Journal of the American Board of Family Medicine*, 30(6), 816–823.
6. Borzecki, A. M., Christiansen, C. L., Chew, P., Loveland, S., & Rosen, A. K. (2010). Comparison of in-hospital versus 30-day mortality assessments for selected medical conditions. *Medical care*, 48(12), 1117–1121.